## Source: Quality Health Care: A Guide to Developing and Using Indicators 2<sup>nd</sup> Edition

**Table 4-5** Advantages and disadvantages of various sampling methods

Sampling Method	Description	Advantages	Disadvantages			
Probability Sampling Methods						
Simple random sample	A sample that is drawn random sample in such a way that every member of a population has an equal chance of being included. A random number table or a random number generator is typically used to actually pull the sample.	<ul> <li>Requires minimum knowledge of the population in advance</li> <li>Free of possible classification errors</li> <li>Easy to analyze the data and compute errors</li> <li>Fairly inexpensive</li> </ul>	<ul> <li>Does not take advantage of the knowledge the researcher might have about the population</li> <li>There could be over- or underrepresentation of subgroups within the population</li> <li>Typically produces larger sampling errors for the same sample than a stratified sample</li> </ul>			
Stratified random sample	The population is divided into relevant strata before random sampling is applied to each stratum.	<ul> <li>Helps to reduce the chances of over-/underrepresenting subgroups within the population</li> <li>Allows you to segment the data into "buckets" during the analysis phase</li> <li>Create more efficient samples</li> <li>Reduces sampling error</li> </ul>	<ul> <li>Requires knowledge of the presence of various characteristics within the population</li> <li>Sampling costs can increase if knowledge of the population is shallow</li> <li>If the strata are not highly homogeneous then sampling error goes up and efficiency goes down</li> </ul>			
Proportional stratified random sample	The proportion (or percentage) of a particular stratum is determined in the population and then applied to the random sample.	<ul> <li>Adds even more precision than the stratified random sample</li> <li>Increases sample representativeness</li> <li>Creates very efficient samples</li> <li>Reduces sampling error</li> </ul>	<ul> <li>Requires more human and financial resources than other methods</li> <li>Requires even more information about the population than stratified random methods</li> </ul>			

This table can be found in *Quality Health Care: A Guide to Developing and Using Indicators 2<sup>nd</sup> Edition* by Dr Robert Lloyd, published by Jones & Bartlett Learning, copyright 2019. Replication and/or distribution of this table without expressed written permission form Dr. Lloyd and Jones & Bartlett Learning is strictly prohibited.

Sampling Method	Description	Advantages	Disadvantages
Systematic sample	Select every $k^{\text{th}}$ observation from the population after a random starting point has been selected.	<ul> <li>Very easy to conduct</li> <li>Has "intuitive" appeal</li> <li>Inexpensive to conduct</li> </ul>	<ul> <li>Can produce bias due to periodic ordering of observation, which produces exclusion of segments of the population</li> <li>Increased probability of sampling bias</li> </ul>
Cluster sample	Clusters or "bunches" of the population are identified, and then random sampling is applied to each cluster.	<ul> <li>Can be low cost, especially if geographic clusters are used</li> <li>If properly done, each cluster is a small model of the population</li> <li>High level of practicality</li> </ul>	<ul> <li>Clusters need to be as heterogeneous as possible</li> <li>Typically has lower statistical efficiency</li> <li>Large samples are often needed to ensure precision</li> </ul>
	Nonprobabilit	y Sampling Methods	
	selected based on availability and convenience. Also known as "accidental" samples.	<ul> <li>Ease of obtaining a sample</li> <li>Relatively low cost</li> </ul>	<ul> <li>Extremely low generalizability</li> <li>No way to determine sampling bias or sampling error</li> </ul>
Quota sample	A population is divided into relevant strata. The desired proportion of samples to be obtained from each stratum is determined, and then a fixed quota within each stratum is set.	<ul> <li>Stratification effect is achieved if the strata are appropriately structured</li> <li>In theory, the quota sample should be reasonably representative of the population</li> <li>Human and financial costs can be kept to a minimum if the strata from which the quotas are to be drawn are grouped close together (reduced the amount of travel the data collectors have to perform in order to gather the data)</li> </ul>	<ul> <li>The people assigned to collect the quotas need to be scrupulous, free from selection bias, and follow the prescribed sampling design (otherwise this method becomes a convenience sample)</li> <li>It is difficult to guarantee that the quotas were filled accurately</li> <li>In-depth knowledge of the population is required</li> <li>Nonrandom selection of the quotas can also introduce bias</li> </ul>

This table can be found in *Quality Health Care: A Guide to Developing and Using Indicators 2<sup>nd</sup> Edition* by Dr Robert Lloyd, published by Jones & Bartlett Learning, copyright 2019. Replication and/or distribution of this table without expressed written permission form Dr. Lloyd and Jones & Bartlett Learning is strictly prohibited.

Sampling Method	Description	Advantages	Disadvantages
Judgment sample	Subgroups are drawn	Samples in a	Sampling bias and
	from a process over	subgroup can be small	sampling error cannot
	time based on expect	"(3–5) since many	be calculated
	knowledge. The	subgroups will be	• Expert knowledge of
	subgroup samples can	selected	the process or
	be drawn either by	Data collection costs	population is required
	random or nonrandom	can be reduced	• Generalization of the
	procedures.	• Provides a dynamic	judgment sample to
		picture of the data and	larger populations
		serves as the basis for	cannot be done
		process improvement	<ul> <li>Personal bias enters</li> </ul>
		Minimum	into the selection of
		stratification effect is	the sample
		achieved	

This table can be found in *Quality Health Care: A Guide to Developing and Using Indicators 2<sup>nd</sup> Edition* by Dr Robert Lloyd, published by Jones & Bartlett Learning, copyright 2019. Replication and/or distribution of this table without expressed written permission form Dr. Lloyd and Jones & Bartlett Learning is strictly prohibited.